

First-Order Continuous Induction and a Logical Study of Real Closed Fields

Saeed Salehi¹  · Mohammadsaleh Zarza²

Received: 1 November 2018 / Revised: 1 May 2019 / Accepted: 7 May 2019 / Published online: 24 June 2019
© Iranian Mathematical Society 2019

Abstract

Over the last century, the principle of “continuous induction” has been studied by different authors in different formats. All of these different forms are equivalent to one of the three versions that we isolate in this paper. We show that one of the three forms of continuous induction is weaker than the other two by proving that it is equivalent to the Archimedean property, while the other two stronger versions are equivalent to the completeness property (the supremum principle) of the real numbers. We study some equivalent axiomatizations for the first-order theory of real closed fields and show that some first-order formalization of continuous induction is able to completely axiomatize it (over the theory of ordered fields).

Keywords First-order logic · Complete theories · Axiomatizing the field of real numbers · Continuous induction · Real closed fields

Mathematics Subject Classification 03B25 · 03C35 · 03C10 · 12L05

Communicated by Ali Enayat.

This is a part of the Ph.D. thesis of the second author written under the supervision of the first author at the University of Tabriz, IRAN.

✉ Saeed Salehi
salehipour@tabrizu.ac.ir
Mohammadsaleh Zarza
m_zarza@tabrizu.ac.ir

¹ Research Institute for Fundamental Sciences, University of Tabriz, 29 Bahman Boulevard, P.O.Box 51666-16471, Tabriz, Iran

² Department of Mathematics, University of Tabriz, P.O.Box 51666-16471, Tabriz, Iran

1 Introduction

Real Analysis is more than a haphazard accumulation of facts about the ordered field of real numbers \mathbb{R} . Indeed, real analysis is a systematic study of \mathbb{R} (and functions on \mathbb{R} , etc.). The most usual systematic way of studying mathematical objects is via axiomatizations. Some axiomatic systems are just definitions; such as the axioms of group theory. Indeed, examples of groups abound in mathematics and other scientific fields. Nonetheless, some axiomatizations are much deeper than definitions; one such example is the axiom system of complete ordered fields. An ordered field is *complete* when it contains the supremum (least upper bound) of every nonempty and bounded subset of itself (let us note that this notion of completeness is formalized in second-order logic). There are very many ordered fields in mathematics, but only one of them, up to isomorphism, is complete (by Dedekind's Theorem; see e.g., [14]). The assumption of the existence of a complete ordered field is not a trivial one, though many textbooks on mathematical real analysis start off with the axioms of complete ordered fields and take \mathbb{R} as a (indeed, as *the*) model of this theory. Of course, this is not the only way to do real analysis; by the arithmetization of analysis, we can construct \mathbb{R} from \mathbb{Q} , and \mathbb{Q} from \mathbb{Z} , and finally \mathbb{Z} from \mathbb{N} . It should be pointed out that we can also reverse this foundational setup by starting with the assumption that \mathbb{R} is a complete ordered field; then we can construct \mathbb{Q} as the smallest subfield of \mathbb{R} , \mathbb{Z} as the smallest sub-ring of \mathbb{R} that contains 1, and \mathbb{N} as the non-negative elements of \mathbb{Z} .

Apart from these philosophical and foundational concerns, here we are interested in the properties of the real numbers \mathbb{R} as a complete ordered field. Indeed, there are several different axiomatizations for the (second-order) theory of complete ordered fields, over the theory of ordered fields **OF**: (1) the existence of supremum (the least upper bound) for every nonempty and bounded subset; (2) the existence of infimum (the greatest lower bound) for every nonempty and bounded subset; and (3) the nonexistence of cuts (with gaps), i.e., a partition into two disjoint subsets in such a way that every element of the first set is smaller than all the elements of the second set, and the first set has no greatest element and the second set has no smallest element. Interestingly, there are many different equivalent statements for the completeness axiom; indeed, as many as 72 of them are listed in [5].

As shown by Tarski, the first-order theory of \mathbb{R} , i.e., the set of first-order sentences in the language of ordered fields $\mathcal{L}_{\mathbf{OF}} = \{+, 0, -, <, \times, 1\}$ that hold in the ordered field of real numbers is axiomatizable by a computable set of axioms known to modern algebraists as **RCF** (real closed fields—see Definition 4.1). Therefore, **RCF** is a complete theory, i.e., any first-order sentence formulated in the language of ordered fields is either provable or refutable in **RCF**. This is in sharp contrast to the semiring \mathbb{N} of natural numbers, the ring \mathbb{Z} of integers, and the field \mathbb{Q} of rationals, none of whose first-order (complete) theories can be axiomatized by a computable set of axioms, thanks to the results of Kurt Gödel (for \mathbb{N} and \mathbb{Z}) and Julia Robinson (for \mathbb{Q}). Thus, the formalization of the fundamentals of Real Analysis naturally leads to central concepts in modern algebra. As we shall see below, formalizing various completeness axioms for the ordered field of real numbers may give rise to various axiomatizations of the theory of real closed fields.

In Sect. 2, we isolate three schemes of the principle of continuous induction, since all the different formats of this principle that have appeared in the literature over the last century are equivalent to one of these three versions. We will show that two of them are equivalent to each other, and to the completeness principle over **OF**, while the third one is weaker. We formalize these three schemes in first-order logic and will compare their strength with each other. In Sect. 3 we study some first-order formalizations of the completeness axiom (of ordered fields) and will see their equivalence with one another by first-order proofs. We will show that the weak principle of continuous induction is equivalent to the Archimedean property (of ordered abelian groups), and the first-order formalization of the (two equivalent) strong principle(s) of continuous induction can axiomatize the theory of real closed fields (over **OF**). In Sect. 4, we study the theory of real closed fields more deeply and introduce a first-order scheme of the Fundamental Theorem of Algebra as an alternative axiomatization of this theory over **OF**. In Sect. 5, we summarize the new and old results of the paper and propose a set of open problems for future investigations. In the Appendix we present a slightly modified proof of Kreisel and Krivine [11, Chapter 4, Theorem 7] for Tarski’s Theorem on the completeness (and the decidability of a first-order axiomatization) of the theory of real closed fields.

2 Continuous Induction, Formalized in First-Order Logic

“Continuous induction”, “induction over the continuum”, “real induction”, “non-discrete induction”, or the like, are some terms used by authors for referring to some statements about the continuum \mathbb{R} . These statements are as strong as the Completeness Axiom of \mathbb{R} and a motivation for their introduction into the literature of mathematics is the easy and sometimes unified ways they provide for proving some basic theorems of mathematical analysis. Here, we do not intend to give a thorough history of the subject or list all of the relevant literature in the References. For a “telegraphic history” we refer the reader to [10], and for an introduction to the subject we refer to [4] and the references therein. The earliest use of continuous induction is perhaps in the paper [3] dating back to 1919. Below, we will formalize it in first-order logic. We will also formalize the formulations of continuous induction presented in [4, 8–10] and later will compare their strength with each other. Let us finally note that [12] is the only textbook (in Persian/Farsi) in which we could find some mention of continuous induction (referring to [9]).

The principle of continuous induction introduced in [3] (see also [4]) is equivalent to the following statement:

Definition 2.1 (**CI**₁) For any $S \subseteq \mathbb{R}$, if

1. for some $a \in \mathbb{R}$ we have $] -\infty, a] \subseteq S$, and
2. there exists some $\epsilon > 0$ such that for all $x \in \mathbb{R}$ if $x \in S$, then $[x, x + \epsilon] \subseteq S$,

then $S = \mathbb{R}$. □

Its formalization in first-order logic on a language \mathcal{L} , where $\{<, 0, +\} \subseteq \mathcal{L}$, is as:

Definition 2.2 (**DCI**₁) Let **DCI**₁(\mathcal{L}), definable continuous induction on \mathcal{L} , be the following first-order scheme

$$\exists x \forall y \leq x \varphi(y) \wedge \exists \epsilon > 0 \forall x (\varphi(x) \rightarrow \forall y [x \leq y \leq x + \epsilon \rightarrow \varphi(y)]) \longrightarrow \forall x \varphi(x),$$

where φ is an arbitrary \mathcal{L} -formula. \square

Continuous induction in [10] (see also [4,9]) is equivalent to the following:

Definition 2.3 (**CI**) For any $S \subseteq \mathbb{R}$, if

1. for some $a \in \mathbb{R}$ we have $] - \infty, a[\subseteq S$, and
2. for any $x \in \mathbb{R}$ if $] - \infty, x[\subseteq S$, then there exists some $\epsilon > 0$ such that $] - \infty, x + \epsilon[\subseteq S$,

then $S = \mathbb{R}$. \square

Remark 2.4 Let us note that this form of real induction can be formulated by using the order relation $<$ only:

For any $S \subseteq \mathbb{R}$, if

1. for some $a \in \mathbb{R}$ we have $] - \infty, a[\subseteq S$, and
2. for any $x \in \mathbb{R}$ if $] - \infty, x[\subseteq S$, then there exists some $y > x$ such that $] - \infty, y[\subseteq S$,

then $S = \mathbb{R}$. \square

The first-order formalization of **CI** is as follows:

Definition 2.5 (**DCI**) Let **DCI**(\mathcal{L}), for a first-order language \mathcal{L} that contains $<$, be the first-order scheme

$$\exists x \forall y < x \varphi(y) \wedge \forall x [\forall y < x \varphi(y) \rightarrow \exists z > x \forall y < z \varphi(y)] \longrightarrow \forall x \varphi(x)$$

for an arbitrary \mathcal{L} -formula φ . \square

When the first-order language \mathcal{L} contains 0 and $+$ also, then this can be formalized as the following:

$$\exists x \forall y < x \varphi(y) \wedge \forall x [\forall y < x \varphi(y) \rightarrow \exists \epsilon > 0 \forall y < x + \epsilon \varphi(y)] \longrightarrow \forall x \varphi(x).$$

Note that in **DCI**₁(\mathcal{L}) there exists a fixed $\epsilon > 0$ such that for all x the second assumption holds, but in **DCI**(\mathcal{L}) for any x there exists some $\epsilon_x > 0$ (which depends on x) such that the second assumption holds.

Finally, there exists a third version of continuous induction which appears in [8]:

Definition 2.6 (**CI**₂) For any $S \subseteq \mathbb{R}$, if

1. for some $a \in \mathbb{R}$ we have $] - \infty, a] \subseteq S$, and
2. for any $x \in \mathbb{R}$ if $(-\infty, x] \subseteq S$, there exists some $y > x$ such that $] - \infty, y[\subseteq S$, and
3. for any $x \in \mathbb{R}$ if $] - \infty, x[\subseteq S$, then $x \in S$,

then $S = \mathbb{R}$. \square

Its formalization in the first-order languages \mathcal{L} that contain $<$ is as follows:

Definition 2.7 (\mathbf{DCI}_2) Let $\mathbf{DCI}_2(\mathcal{L})$ denote the first-order scheme

$$\exists x \forall y \leq x \varphi(y) \wedge \forall x [\forall y \leq x \varphi(y) \rightarrow \exists z > x \forall y < z \varphi(y)] \wedge \forall x [\forall y < x \varphi(y) \rightarrow \varphi(x)] \rightarrow \forall x \varphi(x)$$

where φ is an arbitrary \mathcal{L} -formula. □

Now, we compare the strength of these three schemes with each other.

Theorem 2.8 ($\mathbf{DCI} \iff \mathbf{DCI}_2$) *In any linearly ordered structure $\langle D; \mathcal{L} \rangle$, where \mathcal{L} contains $<$, the scheme $\mathbf{DCI}(\mathcal{L})$ holds, if and only if $\mathbf{DCI}_2(\mathcal{L})$ holds.*

Proof The proof of $\mathbf{DCI}(\mathcal{L}) \implies \mathbf{DCI}_2(\mathcal{L})$ is easy and left to the reader. For $\mathbf{DCI}_2(\mathcal{L}) \implies \mathbf{DCI}(\mathcal{L})$, suppose that $\mathbf{DCI}_2(\mathcal{L})$ holds in a linearly ordered structure $\langle D; \mathcal{L} \rangle$ and assume that for an \mathcal{L} -formula $\varphi(x)$ and some $a \in D$ we have

- (i) $\forall y < a \varphi(y)$, and
- (ii) $\forall x [\forall y < x \varphi(y) \rightarrow \exists z > x \forall y < z \varphi(y)]$.

Now, we show that the following relations hold:

- (1) $\exists x \forall y \leq x \varphi(y)$,
- (2) $\forall x [\forall y \leq x \varphi(y) \rightarrow \exists z > x \forall y < z \varphi(y)]$, and
- (3) $\forall x [\forall y < x \varphi(y) \rightarrow \varphi(x)]$.

This will show, by using $\mathbf{DCI}_2(\mathcal{L})$, that $\forall x \varphi(x)$ holds.

The relation (2) follows straightforwardly from (ii). For (3) fix some $d \in D$ and assume that $\forall y < d \varphi(y)$ holds. Then by (ii), there exists some $d' > d$ such that the statement $\forall y < d' \varphi(y)$ holds too. Whence, $\varphi(d)$ holds as well. The relation (1) holds for $x=a$ for exactly the same reason. □

It follows from the proof of Theorem 2.8 that \mathbf{CI} and \mathbf{CI}_2 are equivalent to each other, in any linear order. Now we show that \mathbf{CI}_1 is strictly weaker than \mathbf{CI} (and \mathbf{CI}_2).

Theorem 2.9 ($\mathbf{DCI} \not\iff \mathbf{DCI}_1$) *For $\mathcal{L} \supseteq \{+, 0, <\}$, in any \mathcal{L} -structure of the form $\langle D; +, 0, <, \dots \rangle$, where $\langle D; +, 0, < \rangle$ is an ordered divisible abelian group, if $\mathbf{DCI}(\mathcal{L})$ holds, then $\mathbf{DCI}_1(\mathcal{L})$ holds too.*

The converse is not true: for $\mathcal{L}_{\mathbf{OF}} = \{+, 0, -, <, \times, 1\}$ the language of ordered fields, the scheme $\mathbf{DCI}_1(\mathcal{L}_{\mathbf{OF}})$ holds in $\langle \mathbb{Q}; \mathcal{L}_{\mathbf{OF}} \rangle$, but $\mathbf{DCI}(\mathcal{L}_{\mathbf{OF}})$ does not.

Proof Suppose that $\mathbf{DCI}(\mathcal{L})$ holds in such a structure $\langle D; +, 0, <, \dots \rangle$. For proving the scheme $\mathbf{DCI}_1(\mathcal{L})$, suppose that for some \mathcal{L} -formula φ and some $\epsilon, a \in D$ with $\epsilon > 0$, we have (i) $\forall y \leq a \varphi(y)$ and (ii) $\forall x (\varphi(x) \rightarrow \forall y [x \leq y \leq x + \epsilon \rightarrow \varphi(y)])$. Then, (1) $\forall y < a \varphi(y)$ holds and we show that (2) $\forall x [\forall y < x \varphi(y) \rightarrow \exists z > x \forall y < z \varphi(y)]$ holds too: for some fixed $b \in D$ assume that $\forall y < b \varphi(y)$. Then we have $\varphi(b - \frac{\epsilon}{2})$ and so (ii) implies that $\forall y [b - \frac{\epsilon}{2} \leq y \leq b + \frac{\epsilon}{2} \rightarrow \varphi(y)]$. Whence, $\forall y < b + \frac{\epsilon}{2} \varphi(y)$ holds, which proves (2). So, $\mathbf{DCI}(\mathcal{L})$ implies $\forall x \varphi(x)$ from (1), (2).

Now, we show that $\mathbf{DCI}_1(\mathcal{L}_{\mathbf{OF}})$ holds in the rational numbers \mathbb{Q} , but $\mathbf{DCI}(\mathcal{L}_{\mathbf{OF}})$ does not. For proving $\mathbb{Q} \models \mathbf{DCI}_1(\mathcal{L}_{\mathbf{OF}})$, suppose that for some φ and $\epsilon, r \in \mathbb{Q}$ with $\epsilon > 0$, we have (i) $\forall y \leq r \varphi(y)$ and (ii) $\forall x (\varphi(x) \rightarrow \forall y [x \leq y \leq x + \epsilon \rightarrow \varphi(y)])$. We show that $\forall x \varphi(x)$ holds in \mathbb{Q} ; if not, then $\mathcal{A} = \{q \in \mathbb{Q} \mid \neg \varphi(q)\}$ is nonempty and bounded below by r . Let $\alpha = \inf \mathcal{A} (\in \mathbb{R})$. So, there exists some $s \in \mathcal{A}$ with $\alpha < s < \alpha + \frac{\epsilon}{2}$, and there exists some $t \in \mathbb{Q}$ with $\alpha - \frac{\epsilon}{2} < t < \alpha$. By $t < \alpha = \inf \mathcal{A}$, we have $t \notin \mathcal{A}$ and so $\varphi(t)$ holds. Then by (ii), we should have $\varphi(y)$ for all $y \in [t, t + \epsilon]$, and in particular $\varphi(s)$, since we already have $t < \alpha < s < \alpha + \frac{\epsilon}{2} < t + \epsilon$; which is a contradiction with $s \in \mathcal{A}$. We now show that $\mathbf{DCI}(\mathcal{L}_{\mathbf{OF}})$ does not hold in $\langle \mathbb{Q}; \mathcal{L}_{\mathbf{OF}} \rangle$. Let $\varphi(x) = [x < 0 \vee x^2 < 2]$. Obviously, (1) $\forall y < 0 \varphi(y)$ holds and in the following we show that (2) $\forall x [\forall y < x \varphi(y) \rightarrow \exists \epsilon > 0 \forall y < x + \epsilon \varphi(y)]$ holds as well. But clearly, $\forall x \varphi(x)$ does not hold since $\varphi(2)$ is not true; this will show that $\mathbf{DCI}(\mathcal{L}_{\mathbf{OF}})$ is not true in \mathbb{Q} . For showing (2), fix an $r \in \mathbb{Q}$ and assume that $\forall y < r \varphi(y)$. We now prove that $\exists \epsilon > 0 \forall y < r + \epsilon \varphi(y)$.

- (I) If $r < 0$, then for $\epsilon = -\frac{r}{2}$ we have $\forall y < r + \epsilon \varphi(y)$.
 (II) If $r = 0$, then for $\epsilon = 1$ we have $\forall y < r + \epsilon \varphi(y)$.
 (III) If $r > 0$, then we have $r^2 < 2$, since $r^2 = 2$ is impossible (by $\sqrt{2} \notin \mathbb{Q}$) and if $r^2 > 2$, then we cannot have $\forall y < r \varphi(y)$ because for $y = \frac{r^2+2}{2r}$ we have $0 < y < r$ and $y^2 - 2 = (\frac{r^2-2}{2r})^2 > 0$. Let $\epsilon = \min\{1, \frac{2-r^2}{2r+1}\}$, then $0 < \epsilon \leq 1$. For showing $\forall y < r + \epsilon \varphi(y)$, take some y with $0 \leq y < r + \epsilon$. We have $y^2 - 2 < (r + \epsilon)^2 - 2 = (r^2 - 2) + \epsilon(2r + \epsilon) \leq (r^2 - 2) + \epsilon(2r + 1) \leq (r^2 - 2) + (2 - r^2) = 0$. So, $y^2 < 2$, thus $\varphi(y)$ holds for any $y < r + \epsilon$. \square

Therefore, \mathbf{CI}_1 cannot be regarded as a genuine “continuous induction”, even though it is a kind of “non-discrete induction” since it does not hold in \mathbb{Z} or \mathbb{N} . But since it holds in \mathbb{Q} , it is not really an “induction on the continuum”. In the next section we will see the real reason for regarding \mathbf{CI} (and also its equivalent \mathbf{CI}_2) as a genuine “continuous induction”, and not regarding \mathbf{CI}_1 as a true “induction on the continuum”.

3 First-Order Completeness Axioms for the Field of Real Numbers

In mathematical analysis, \mathbb{R} is usually introduced as a complete ordered field; indeed by a theorem of Dedekind there exists only one complete ordered field up to isomorphism. In most of the textbooks on mathematical analysis, the existence of a complete ordered field is assumed as an axiom; in some other textbooks, a complete ordered field is constructed from \mathbb{Q} either by Dedekind cuts or by Cauchy sequences. The most usual completeness axiom in the analysis textbooks is the existence of supremum for any non-empty and bounded above subset; its first-order version is as follows (we assume that \mathcal{L} always contains $<$):

Definition 3.1 (D-Sup) Let $\mathbf{D-Sup}(\mathcal{L})$, Definable Supremum Property, be the following first-order scheme

$$\exists x \varphi(x) \wedge \exists y \forall x [\varphi(x) \rightarrow x \leq y] \longrightarrow \exists z \forall y (\forall x [\varphi(x) \rightarrow x \leq y] \leftrightarrow z \leq y),$$

where φ is an arbitrary \mathcal{L} -formula. □

Informally, **D-Sup** says that for a nonempty $(\exists x \varphi(x))$ and bounded from above set (for some y we have $\forall x [\varphi(x) \rightarrow x \leq y]$) there exists a least upper bound (for some z , any y is an upper bound if and only if $y \geq z$). A dual statement is the principle of the existence of infimum for any nonempty and bounded from below set:

Definition 3.2 (D-Inf) Let **D-Inf**(\mathcal{L}) be the first-order scheme

$$\exists x \varphi(x) \wedge \exists y \forall x [\varphi(x) \rightarrow y \leq x] \longrightarrow \exists z \forall y (\forall x [\varphi(x) \rightarrow y \leq x] \leftrightarrow y \leq z)$$

for arbitrary \mathcal{L} -formula φ . □

Indeed, the scheme **D-Inf** is used by Tarski for presenting a complete first-order axiomatic system for the ordered field of real numbers (see [6]). The usual real analytic proof for the equivalence of these two schemes works in first-order logic as well:

Theorem 3.3 (D-Sup \iff D-Inf) *In any linearly ordered structure $\langle D; \mathcal{L} \rangle$, **D-Sup**(\mathcal{L}) holds if and only if **D-Inf**(\mathcal{L}) holds.*

Proof We prove only **D-Sup**(\mathcal{L}) \implies **D-Inf**(\mathcal{L}); its converse can be proved by a dual argument. Suppose that for some $\varphi(x)$ and $a, b \in D$ we have (i) $\varphi(a)$ and (ii) $\forall x [\varphi(x) \rightarrow b \leq x]$. For using **D-Sup**(\mathcal{L}), let $\psi(x) = \forall y < x \neg \varphi(y)$. Then, by (ii), we have (1) $\psi(b)$ and by (i) we have (2) $\forall x [\psi(x) \rightarrow x \leq a]$. Now, by **D-Sup**(\mathcal{L}) there exists some $c \in D$ such that

$$\forall y (\forall x [\psi(x) \rightarrow x \leq y] \leftrightarrow c \leq y). \tag{3.1}$$

We show $\forall z (\forall x [\varphi(x) \rightarrow z \leq x] \leftrightarrow z \leq c)$. For every $z \in D$ we have:

- (I) If $\forall x [\varphi(x) \rightarrow z \leq x]$, then $\psi(z)$; from (3.1) we have $\forall x [\psi(x) \rightarrow x \leq c]$ so $z \leq c$.
- (II) If $z \leq c$, then for any $x < z$ we have $c \not\leq x$ and so (3.1) implies that for some u we have $\psi(u)$ and $u \not\leq x$. Now, $\psi(u)$ and $x < u$ imply that $\neg \varphi(x)$. Thus, $\forall x < z \neg \varphi(x)$, or equivalently, $\forall x [\varphi(x) \rightarrow z \leq x]$. □

Another completeness principle is the so called Dedekind’s Axiom; which says that there is no proper cut (with a gap) in \mathbb{R} . The following is a first-order writing of an equivalent form of this axiom:

Definition 3.4 (D-Cut) Let **D-Cut**(\mathcal{L}) denote the first-order scheme

$$\begin{aligned} &\exists x \exists y [\varphi(x) \wedge \psi(y)] \wedge \forall x \forall y [\varphi(x) \wedge \psi(y) \rightarrow x < y] \\ &\longrightarrow \exists z \forall x \forall y [\varphi(x) \wedge \psi(y) \rightarrow x \leq z \leq y], \end{aligned}$$

where φ and ψ are arbitrary \mathcal{L} -formulas. □

Tarski used this scheme for presenting another complete first-order axiomatic system for \mathbb{R} (see [6]). Indeed, in linearly ordered \mathcal{L} -structures, **D-Cut**(\mathcal{L}) is equivalent to the other completeness axioms schemes of **D-Sup**(\mathcal{L}) and **D-Inf**(\mathcal{L}):

Theorem 3.5 ($\mathbf{D-Sup} \iff \mathbf{D-Cut}$) *In any linearly ordered structure $\langle D; \mathcal{L} \rangle$, $\mathbf{D-Sup}(\mathcal{L})$ holds if and only if $\mathbf{D-Cut}(\mathcal{L})$ holds.*

Proof If $\mathbf{D-Sup}(\mathcal{L})$ holds in a linear order $\langle D; \mathcal{L} \rangle$, then for showing $\mathbf{D-Cut}(\mathcal{L})$ assume that for formulas $\varphi(x)$ and $\psi(x)$ and elements $a, b \in D$ we have (i) $\varphi(a) \wedge \psi(b)$ and (ii) $\forall x \forall y [\varphi(x) \wedge \psi(y) \rightarrow x < y]$; we show the existence of some element $z \in D$ such that $\forall y [\varphi(x) \wedge \psi(y) \rightarrow x \leq z \leq y]$. Now, by (i), we have (1) $\varphi(a)$ and, by (ii), we have (2) $\forall x [\varphi(x) \rightarrow x < b]$. By $\mathbf{D-Sup}(\mathcal{L})$ there exists some element $c \in D$ such that (3) $\forall y (\forall x [\varphi(x) \rightarrow x \leq y] \iff c \leq y)$. We show that $\forall x \forall y [\varphi(x) \wedge \psi(y) \rightarrow x \leq c \leq y]$ holds. By (3), for $y = c$, we have $x \leq c$ for any x with $\varphi(x)$. Now assume that $\psi(y)$; then if $c \not\leq y$, by (3), there should exist some x such that $\varphi(x)$ but $y < x$. This contradicts (ii), since on the one hand we have $\varphi(x) \wedge \psi(y)$, but on the other hand $y < x$!

If $\mathbf{D-Cut}(\mathcal{L})$ holds in a linearly ordered $\langle D; \mathcal{L} \rangle$, then for showing $\mathbf{D-Sup}(\mathcal{L})$ assume that for some $\varphi(x)$ and $a, b \in D$ we have (i) $\varphi(a)$ and (ii) $\forall x [\varphi(x) \rightarrow x \leq b]$; we show the existence of some $z \in D$ such that $\forall y (\forall x [\varphi(x) \rightarrow x \leq y] \iff z \leq y)$. We distinguish two cases:

(I) For some $d \in D$ we have $\varphi(d) \wedge \forall x [\varphi(x) \rightarrow x \leq d]$. In this case, it is easy to show that $\forall y (\forall x [\varphi(x) \rightarrow x \leq y] \iff d \leq y)$ holds, since for any y we have $y < d$ if and only if for some x (which is $x = d$) we have $x > y$ and $\varphi(x)$.

(II) We have $\forall y [\forall x [\varphi(x) \rightarrow x \leq y] \rightarrow \neg \varphi(y)]$. In this case, put $\psi(x)$ be the formula $\psi(x) = \forall y \geq x \neg \varphi(y)$. Then, noting that by (ii) we already have $\neg \varphi(b)$, the relation $\psi(b)$ holds, and so by (i) we have (1) $\varphi(a) \wedge \psi(b)$. Also, (2) $\forall x \forall y [\varphi(x) \wedge \psi(y) \rightarrow x < y]$ holds by the definition of ψ . Thus, by $\mathbf{D-Cut}(\mathcal{L})$, there exists some $c \in D$ such that (3) $\forall x \forall y [\varphi(x) \wedge \psi(y) \rightarrow x \leq c \leq y]$. We show that this c satisfies $\forall y (\forall x [\varphi(x) \rightarrow x \leq y] \iff c \leq y)$.

First, assume that $\forall x [\varphi(x) \rightarrow x \leq y]$; then $\neg \varphi(y)$ holds by the assumption (II). So, $\psi(y)$ holds and therefore $c \leq y$ by (3).

Second, assume that $c \leq y$ but $\varphi(x) \wedge x \not\leq y$ for some x . Then by (3), and $\varphi(x)$, we should have $x \leq c$, but this contradicts $c \leq y < x$! \square

Finally, we observe that continuous induction is equivalent to (any of) the (above) completeness axiom(s), even in first-order logic:

Theorem 3.6 ($\mathbf{D-Inf} \iff \mathbf{DCI}$) *In any linearly ordered structure $\langle D; \mathcal{L} \rangle$, $\mathbf{D-Inf}(\mathcal{L})$ holds if and only if $\mathbf{DCI}(\mathcal{L})$ holds.*

Proof Suppose that $\mathbf{D-Inf}(\mathcal{L})$ holds in a linearly ordered structure $\langle D; \mathcal{L} \rangle$. For proving $\mathbf{DCI}(\mathcal{L})$, assume that, for some $\varphi(x)$ and $a \in D$, we have (i) $\forall y < a \varphi(y)$ and (ii) $\forall x [\forall y < x \varphi(y) \rightarrow \exists z > x \forall y < z \varphi(y)]$; then we show that $\forall x \varphi(x)$. If for some $b \in D$ we had $\neg \varphi(b)$, then put $\varphi'(x) = \neg \varphi(x)$. Now we have (1) $\varphi'(b)$ and (2) $\forall x [\varphi'(x) \rightarrow a \leq x]$ by (i). By the assumption $\mathbf{D-Inf}(\mathcal{L})$, there exists some element $c \in D$ for which we have $\forall y (\forall x [\varphi'(x) \rightarrow y \leq x] \iff y \leq c)$; or, equivalently, (3) $\forall y [y \leq c \iff \forall x < y \varphi(x)]$. Thus, $\forall x < c \varphi(x)$ and so by (ii) there exists some $d \in D$ such that $d > c$ and $\forall y < d \varphi(y)$. By (3), since $d > c$, there exists some $x < d$ such that $\neg \varphi(x)$. This is a contradiction, since we had $\forall y < d \varphi(y)$.

Now, suppose that $\mathbf{DCI}(\mathcal{L})$ holds in such a $\langle D; \mathcal{L} \rangle$; for proving $\mathbf{D-Inf}(\mathcal{L})$ assume that we have (i) $\varphi(a)$ and (ii) $\forall x [\varphi(x) \rightarrow b \leq x]$ for some formula φ

and some $a, b \in D$. We aim at showing the existence of some $z \in D$ such that (iii) $\forall y (\forall x [\varphi(x) \rightarrow y \leq x] \leftrightarrow y \leq z)$. Assume that for no z the condition (iii) holds; so for any z there exists some z' such that (iv) $\neg[z' \leq z \leftrightarrow \forall x < z' \neg \varphi(x)]$. Now, for $\varphi'(x) = \neg \varphi(x)$ we have (1) $\forall x < b \varphi'(x)$ by (ii), and we show that the condition (2) $\forall x [\forall y < x \varphi'(y) \rightarrow \exists z > x \forall y < z \varphi'(y)]$ holds: for any fixed x_0 if $\forall y < x_0 \varphi'(y)$, then (3) $\forall x < x_0 \neg \varphi(x)$. There exists some z_0 such that (iv) holds (for $z' = z_0$); if $z_0 \leq x_0$, then $\neg \forall x < z_0 \neg \varphi(x)$ or $\exists x < z_0 \varphi(x)$ which contradicts (3). So, we have $x_0 < z_0$; by (iv) we should have $\forall x < z_0 \varphi'(x)$. Whence, (2) holds, and we can apply **DCI**(\mathcal{L}), which implies that $\forall x \varphi'(x)$ and this contradicts (i). Therefore, for some $z \in D$, the statement (iii) should hold. \square

It is proved in [10, Theorems 3.1,3.2] that continuous induction **CI** is equivalent to the infimum (greatest lower bound) property in ordered fields; also, the scheme **CI** is proved in [9, Theorem 1] to be equivalent to Dedekind’s Axiom (the nonexistence of cuts with gaps). We now show that the principle of continuous induction as presented in Definition 2.1 (**CI**₁) is actually equivalent to the Archimedean property in ordered abelian groups.

Definition 3.7 (Archimedean property, **AP**) An ordered abelian group $\langle G; +, 0, < \rangle$ has the Archimedean property when for any $a, b \in G$ with $a > 0$ there exists some $n \in \mathbb{N}$ such that $b < n \cdot a$, where $n \cdot a = \underbrace{a + \dots + a}_{n\text{-times}}$. \square

Theorem 3.8 (**AP** \iff **CI**₁) An ordered abelian group has the Archimedean property if and only if it satisfies **CI**₁.

Proof Suppose that the ordered abelian group $\langle G; +, 0, < \rangle$ has the Archimedean property (**AP**) and we have (i) $] - \infty, a] \subseteq \mathcal{A}$ and (ii) for any $x \in G$, if $x \in \mathcal{A}$, then $[x, x + \epsilon] \subseteq \mathcal{A}$; for a subset $\mathcal{A} \subseteq G$ and some $\epsilon, a \in G$ with $\epsilon > 0$. By induction on $n \in \mathbb{N}$, we can show that (iii) $] - \infty, a + n \cdot \epsilon] \subseteq \mathcal{A}$: for $n = 0$ it follows from (i) and the induction step follows from (ii). For an arbitrary $x \in G$, by **AP** there exists some $n \in \mathbb{N}$ such that $x - a < n \cdot \epsilon$; so we have $x \in \mathcal{A}$ by (iii). Whence, $\mathcal{A} = G$, and so **CI**₁ holds in G .

Now, suppose that the ordered abelian group $\langle G; +, 0, < \rangle$ satisfies **CI**₁. For any $a, b \in G$ with $a > 0$, let $\mathcal{B} = \{x \in G \mid \exists n \in \mathbb{N} : x < n \cdot a\}$. Then we obviously have (i) $] - \infty, 0] \subseteq \mathcal{B}$; we show that (ii) for any $x \in G$, if $x \in \mathcal{B}$, then $[x, x + a] \subseteq \mathcal{B}$. For any $x \in \mathcal{B}$ we have $x < m \cdot a$ for some $m \in \mathbb{N}$, so for any $y \in [x, x + a]$ we have $y \leq x + a < (m + 1) \cdot a$, and so $y \in \mathcal{B}$. Thus, by **CI**₁, from (i) and (ii), we should have $\mathcal{B} = G$, and so for any $b \in G$ there is some $n \in \mathbb{N}$ such that $b < n \cdot a$; whence G has **AP**. \square

Let us note that Theorem 3.8 gives another proof for $\mathbb{Q} \models \mathbf{CI}_1$ (and also $\mathbb{Q} \models \mathbf{DCI}_1$) that was proved already in Theorem 2.9. So, **CI**₁ can be added to the list of 42 equivalent forms of the Archimedean property (of ordered fields) in [5]. Whence, while the principle of continuous induction in Definitions 2.3 (**CI**) and 2.6 (**CI**₂) are equivalent to the completeness axiom (in the ordered fields), the principle **CI**₁ is equivalent to the Archimedean property (and is not equivalent to the completeness axiom) in such fields.

4 Real Closed Fields, a First-Order Logical Study

What makes the real closed fields interesting in mathematical logic is Tarski's theorem that, $\text{Th}(\langle \mathbb{R}; +, 0, -, <, \times, 1 \rangle)$, the complete first-order theory of the ordered field of real numbers is decidable and axiomatizable by the theory of real closed (ordered) fields (see e.g., [11]). The real closed fields are studied thoroughly in algebra and algebraic geometry (see e.g., [1]); here we study some equivalent axiomatizations of the first-order theory of real closed fields. The most usual definition of a real closed field is an ordered field which satisfies the following axiom scheme:

Definition 4.1 (RCF) Let **RCF** be the following axiom scheme:

$$\forall x > 0 \exists y (y^2 = x) \wedge \forall \{a_i\}_{i \leq 2n} \exists x \left(x^{2n+1} + \sum_{i \leq 2n} a_i x^i = 0 \right),$$

for $n \in \mathbb{N}$ with $n > 1$, together with the axioms of Ordered Fields (**OF**). \square

By definition, a real closed field is an ordered field in which every positive element has a square root and every odd-degree polynomial has a root in it. This most usual definition of real closed fields is in fact the most inapplicable one, since many interesting theorems on and about real closed fields use other equivalent definitions. The most applicable (and the most fruitful) definition of real closed fields is the one which says that a real closed field is an ordered field in which the Intermediate Value Theorem (IVT) holds:

Definition 4.2 (IVT) Let **IVT** be the first-order scheme

$$\forall p \forall u, v \exists x [u < v \wedge p(u) \cdot p(v) < 0 \longrightarrow u < x < v \wedge p(x) = 0],$$

where $\forall p$ stands for $\forall \{a_i\}_{i \leq m}$ for polynomial $p(x) = \sum_{i \leq m} a_i x^i$. \square

Let us note that the terms of $\mathcal{L}_{\text{OF}} = \{+, 0, -, <, \times, 1\}$ the language of ordered fields which contain a variable x are polynomials like $p(x) = \sum_{i \leq m} a_i x^i$ where a_i 's are some x -free terms. For the usefulness of **IVT**, we first note that it implies **RCF**:

Theorem 4.3 (IVT \implies RCF) In any ordered field that **IVT** holds, **RCF** holds too.

Proof For any $a > 0$, let $p(x) = x^2 - a$. Then we have $p(0) = -a < 0 < a^2 + \frac{1}{4} = p(a + \frac{1}{2})$, and so by **IVT** for some x with $0 < x < a + \frac{1}{2}$ we have $p(x) = 0$; thus, $x^2 = a$.

We can write any odd-degree polynomial as $p(x) = x^{2n+1} + \sum_{i \leq 2n} a_i x^i$ by multiplying with an x -free term, if necessary. Let $v = 1 + \sum_{i \leq 2n} |a_i|$ and $u = -v$. Then $u \leq -1 < 0 < 1 \leq v$, so $|u|, |v| \geq 1$. Now, by $u + \sum_i |a_i| = -1$ and the triangle inequality, we have

$$p(u) \leq u^{2n+1} + \sum_{i \leq 2n} |a_i| |u|^i \leq u^{2n+1} + \sum_{i \leq 2n} |a_i| u^{2n} = u^{2n} (u + \sum_{i \leq 2n} |a_i|) = -u^{2n} < 0, \text{ and}$$

$$p(v) \geq v^{2n+1} - \sum_{i \leq 2n} |a_i| |v|^i \geq v^{2n+1} - \sum_{i \leq 2n} |a_i| v^{2n} = v^{2n} (v - \sum_{i \leq 2n} |a_i|) = v^{2n} > 0.$$

Finally, the desired conclusion follows from **IVT** by $u < v$ and $p(u) < 0 < p(v)$. \square

As some other uses of the **IVT**, let us take a look at some real analytic theorems that are proved algebraically (cf. [1]).

Definition 4.4 (*Derivative*) The derivative of a polynomial $p(x) = \sum_{i \leq n} a_i x^i$ is the polynomial $p'(x) = \sum_{0 < i \leq n} i a_i x^{i-1}$.

Remark 4.5 (*Some properties of derivatives*) For polynomials $p(x) = \sum_{i \leq n} a_i x^i$ and $q(x) = \sum_{j \leq m} b_j x^j$, we have:

- $(ap)' = ap'$ for a constant (i.e., an x -free term) a ,
- $(p + q)' = p' + q'$, and
- $(p \cdot q)' = (p' \cdot q) + (p \cdot q')$,

which can be verified rather easily. For example, the last item can be verified by noting that the coefficient of x^{k-1} in $(p \cdot q)'$ is $k \sum_{i+j=k} a_i b_j = \sum_{i \leq k} k a_i b_{k-i} = \sum_{i \leq k} [i a_i b_{k-i} + (k-i) a_i b_{k-i}] = [\sum_{i \leq k} (i a_i) b_{k-i}] + [\sum_{j \leq k} a_{k-j} (j b_j)]$ in which the first summand is the coefficient of x^{k-1} in $(p' \cdot q)$ and the second summand is the coefficient of x^{k-1} in $(p \cdot q')$. □

It goes without saying that the properties mentioned in Remark 4.5 are the ones that are usually learned in elementary calculus through the analytic methods (cf. [14]); so are the following theorem and lemma.

Theorem 4.6 (**IVT** \implies **Rolle** + **MVT** + **Derivative Signs**) Let F be an ordered field in which **IVT** holds. Let $p(x)$ be a polynomial with the coefficients in F and let $a, b \in F$ with $a < b$.

Rolle’s Theorem: If $p(a) = p(b) = 0$, then for some $c \in]a, b[$ we have $p'(c) = 0$.

Mean Value Theorem: There exists some $c \in]a, b[$ such that $p'(c) = \frac{p(b) - p(a)}{b - a}$.

Derivative Signs: If $p'(x) > 0$ (respectively, $p'(x) < 0$) for all $x \in]a, b[$, then $p(u) < p(v)$ (respectively, $p(u) > p(v)$) for any $u, v \in F$ with $a < u < v < b$.

Proof For Rolle’s Theorem, we note that since any polynomial can have only a finite number of roots, then we can assume that there is no root of p in the open interval $]a, b[$. Now, by the assumption $p(a) = p(b) = 0$, both $(x - a)$ and $(x - b)$ divide $p(x)$; let m be the greatest number such that $(x - a)^m$ divides $p(x)$ and n be the greatest number such that $(x - b)^n$ divides $p(x)$. Then we can write $p(x) = (x - a)^m (x - b)^n q(x)$ for some polynomial $q(x)$ that has no root in $]a, b[$, and so by **IVT** we have $q(a)q(b) > 0$. Therefore, we have, by Remark 4.5, that $p'(x) = (x - a)^{m-1} (x - b)^{n-1} r(x)$ where $r(x) = m(x - b)q(x) + n(x - a)q(x) + (x - a)(x - b)q'(x)$. Whence, we have the (in)equality $r(a)r(b) = m(a - b)q(a)n(b - a)q(b) = -mn(b - a)^2 q(a)q(b) < 0$, and so **IVT** implies the existence of some $c \in]a, b[$ with $r(c) = 0$; from which $p'(c) = 0$ follows.

The Mean Value Theorem classically follows (rather straightforwardly) from Rolle’s Theorem for the polynomials $q(x) = p(x) - \frac{p(b) - p(a)}{b - a} x$, and $r(x) = q(x) - q(a)$, since we have $q(a) = q(b)$ and so $r(a) = r(b) = 0$; also, $r'(x) = q'(x) = p'(x) - \frac{p(b) - p(a)}{b - a}$.

Derivative Signs easily follows from the Mean Value Theorem: for any such u, v there exists some $w \in]u, v[$ such that $p(v) - p(u) = p'(w) \cdot (v - u)$. Now, if $p'(w) > 0$, then $p(v) > p(u)$, and if $p'(w) < 0$, then $p(v) < p(u)$. \square

The last (and the strongest) witness for the strength of **IVT** is Tarski's theorem that proves that the theory **OF + IVT** is complete; i.e., for any sentence θ in the language of ordered fields, $\mathcal{L}_{\text{OF}} = \{+, 0, -, <, \times, 1\}$, we have either **OF + IVT** $\vdash \theta$ or **OF + IVT** $\vdash \neg\theta$. In the Appendix we present a somewhat modified proof of Kreisel and Krivine [11] for this result (Theorem 5.1). Let us then note this last result implies that every statement that is true in \mathbb{R} is provable from **IVT (+OF)**, since, otherwise its negation would have been provable, and this would contradict its truth in \mathbb{R} . Thus, for example, we have **IVT** \implies **D-Sup**(\mathcal{L}_{OF}) (and also **IVT** \implies **DCI**(\mathcal{L}_{OF}), etc.) over the theory **OF**. Moreover, if an axiom (or an axiom scheme) that is true in \mathbb{R} can prove **IVT**, then it is actually equivalent to **IVT**, and so can be used as another axiomatization of the theory of real closed fields (over **OF**). So, by the following theorem (4.8), all the schemes that we have considered so far (except **DCI**₁) are equivalent to **IVT**. We will need the following lemma for proving **D-Inf**(\mathcal{L}_{OF}) \implies **IVT**.

Lemma 4.7 (Continuity of polynomials) *For any polynomial q and element w , if $q(w) > 0$ (respectively, if $q(w) < 0$), then there exists some $\epsilon > 0$ such that for any $x \in [w - \epsilon, w + \epsilon]$ we have $q(x) > 0$ (respectively, $q(x) < 0$).*

Proof Let $q(x) = \sum_{k \leq m} a_k x^k$, and suppose for $A > 0$ we have $|w| < A$ and $|a_k| < A$ for $k \leq m$. Let $B = A \sum_{k \leq m} \sum_{i < k} \binom{k}{i} A^i$, and choose an $\epsilon > 0$ such that $\epsilon < 1$ and $\epsilon B < \frac{1}{2}|q(w)|$; note that $|q(w)| \neq 0$. Now, for any $x \in [w - \epsilon, w + \epsilon]$ we have $x = w + \delta$ for some $\delta \in [-\epsilon, \epsilon]$. So,

$$\begin{aligned} |q(x) - q(w)| &= \left| \sum_{k \leq m} a_k ([w + \delta]^k - w^k) \right| = \left| \sum_{k \leq m} a_k \sum_{i < k} \binom{k}{i} \delta^{k-i} w^i \right| \\ &\leq \sum_{k \leq m} |a_k| \sum_{i < k} \binom{k}{i} |\delta|^{k-i} |w|^i \leq \sum_{k \leq m} A \sum_{i < k} \binom{k}{i} \epsilon^{k-i} A^i \\ &= \epsilon A \sum_{k \leq m} \sum_{i < k} \binom{k}{i} \epsilon^{k-1-i} A^i \leq \epsilon B < \frac{1}{2}|q(w)|. \end{aligned}$$

Whence, $q(w) - \frac{1}{2}|q(w)| < q(x) < q(w) + \frac{1}{2}|q(w)|$. So, if $q(w) > 0$, then $q(x) > \frac{1}{2}q(w) > 0$, and if $q(w) < 0$, then $q(x) < \frac{1}{2}q(w) < 0$, for any $x \in [w - \epsilon, w + \epsilon]$. \square

Theorem 4.8 (**D-Inf** \implies **IVT**) *Any ordered field that satisfies **D-Inf**(\mathcal{L}_{OF}) satisfies **IVT** too.*

Proof Suppose that for a polynomial p we have $p(u)p(v) < 0$ for some u, v with $u < v$. Let $\varphi(x) = u \leq x \leq v \wedge p(u)p(x) < 0$. Then $\varphi(v)$ and $\forall x [\varphi(x) \rightarrow u \leq x]$. So, by **D-Inf**(\mathcal{L}_{OF}) there exists some w such that (1) $\forall y (\forall x [\varphi(x) \rightarrow y \leq x] \leftrightarrow y \leq w)$. We show that $u \leq w \leq v$ and also $p(w) = 0$. If $w < u$, then by (1) for some x , we should have

$\varphi(x)$ and $x < u$, and this is impossible; so $u \leq w$. Also, if $v < w$, then there exists some w' such that $v < w' < w$, and so by (1) for any x with $\varphi(x)$ we should have $w' \leq x$ and in particular, since $\varphi(v)$, we should have $w' \leq v$; contradiction, thus $w \leq v$. Now, we show that $p(u)p(w) = 0$. If not, i.e., $p(u)p(w) \neq 0$, then we have either (i) $p(u)p(w) < 0$ or (ii) $p(u)p(w) > 0$. In either case by Lemma 4.7, there exists some $\epsilon > 0$ such that either (i') $\forall x \in [w - \epsilon, w + \epsilon]: p(u)p(x) < 0$ or (ii') $\forall x \in [w - \epsilon, w + \epsilon]: p(u)p(x) > 0$. In case (i') we have $\varphi(w - \epsilon)$, but then by (1), for $y = w$, we should have $w \leq w - \epsilon$, a contradiction. In case (ii') by (1), since $w + \epsilon \not\leq w$, there should exist some x such that $\varphi(x)$ and $x < w + \epsilon$. By (1), for $y = w$, we should have $w \leq x$, from $\varphi(x)$. So, $x \in [w - \epsilon, w + \epsilon]$, but then we should have by (ii') that $p(u)p(x) > 0$ contradicting $\varphi(x)$ (which implies $p(u)p(x) < 0$). Thus, $p(u)p(w) = 0$, and so $p(w) = 0$. Let us note that $w \neq u, v$ by $p(u)p(v) < 0$, and so $w \in]u, v[$. \square

Remark 4.9 (**D-Inf** \implies **RCF**) Theorems 4.8 and 4.3 imply that any ordered field that satisfies **D-Inf**(\mathcal{L}_{OF}) also satisfies **RCF**. This could also be proved directly by elementary analysis (using Lemma 4.7): for a given $a > 0$, the infimum x of the (definable) set $\{u \mid u > 0 \wedge u^2 > a\}$ satisfies $x^2 = a$, and the infimum y of the set $\{u \mid (u^{2n+1} + \sum_{i \leq 2n} a_i u^i) > 0\}$ satisfies $y^{2n+1} + \sum_{i \leq 2n} a_i y^i = 0$. \square

So, from Theorem 5.1 we will have

$$\begin{aligned} \mathbf{IVT} &\equiv \mathbf{D-Inf}(\mathcal{L}_{OF}) \equiv \mathbf{D-Sup}(\mathcal{L}_{OF}) \equiv \mathbf{D-Cut}(\mathcal{L}_{OF}) \\ &\equiv \mathbf{DCI}(\mathcal{L}_{OF}) \equiv \mathbf{DCI}_2(\mathcal{L}_{OF}). \end{aligned}$$

As for **RCF**, we have shown only the conditional **IVT** \implies **RCF** in Theorem 4.3. Its converse (**RCF** \implies **IVT**) is usually proved in the literature by first proving the fundamental theorem of algebra, which says that the field $\mathbb{R}(i)$ is algebraically closed, where $F(i)$ is the result of adjoining $i = \sqrt{-1}$ to F (see, e.g., [1, Theorem 2.11]). A field is called algebraically closed when any polynomial with coefficients in it has a root (in that field). As a matter of fact, an equivalent definition for real closed fields in the literature is that “an ordered field F is real closed if and only if the field $F(i)$ is algebraically closed” (thanks to a theorem of Artin and Schreier). An equivalent statement is that “an ordered field is real closed if and only if every positive element has a square root and any polynomial can be factorized into linear and quadratic factors”. This is also an equivalent form of the fundamental theorem of algebra (for \mathbb{R}), for which we propose the following first-order scheme:

Definition 4.10 (**FTA_{RCF}**) Let **FTA_{RCF}** denote the conjunction of $\forall x > 0 \exists y (y^2 = x)$ and the following first-order scheme

$$\forall \{a_i\}_{i < 2n} \exists \{b_j, c_j\}_{j < n} \forall x \left[\left(x^{2n} + \sum_{i < 2n} a_i x^i \right) = \prod_{j < n} (x^2 + b_j x + c_j) \right]$$

for any $n \in \mathbb{N}$ with $n > 1$. \square

So, $\mathbf{FTA}_{\text{RCF}}$ says that any even-degree polynomial can be factorized as a product of some quadratic polynomials. The following theorem is proved in e.g. [1, Theorem 2.11]; here we present a different proof:


Theorem 4.11 ($\mathbf{FTA}_{\text{RCF}} \implies \mathbf{IVT}$) *Any ordered field that satisfies $\mathbf{FTA}_{\text{RCF}}$ satisfies \mathbf{IVT} too.*

Proof For $p(x)$ with degree m , suppose that $p(u) \cdot p(v) < 0$ for some u, v with $u < v$. Put $q(x) = \frac{1}{p(u)}(1+x^2)^m p(u + \frac{v-u}{1+x^2})$. Then $q(x) = x^{2m} + r(x^2)$ for some $r(x)$ whose degree is less than m . By $\mathbf{FTA}_{\text{RCF}}$ we have $q(x) = \prod_{j < m} (x^2 + b_j x + c_j)$ for some $\{b_j, c_j\}_{j < m}$. Now, we have $\prod_{j < m} c_j = q(0) = \frac{p(v)}{p(u)} = \frac{p(u)p(v)}{p(u)^2} < 0$. So, $c_j < 0$ for some j . Assume, $c_0 < 0$. Whence, $b_0^2 - 4c_0 > 0$ and so $b_0^2 - 4c_0 = d^2$ for some d . Thus, $s = \frac{1}{2}(-b_0 + d)$ is a root of $x^2 + b_0 x + c_0$ and so a root of $q(x)$ too. Finally, for $r = u + \frac{v-u}{1+s^2}$ we have, obviously, $u < r < v$ and $p(r) = 0$. \square

So, by Theorems 4.11 ($\mathbf{FTA}_{\text{RCF}} \implies \mathbf{IVT}$) and 4.3 ($\mathbf{IVT} \implies \mathbf{RCF}$), the scheme $\mathbf{FTA}_{\text{RCF}}$ implies that any odd-degree polynomial has a root. So, one can prove, by an induction on the degree of the polynomials that, when $\mathbf{FTA}_{\text{RCF}}$ holds, every polynomial can be written as the product of some linear and quadratic factors. Let us also note that $\mathbf{FTA}_{\text{RCF}} \implies \mathbf{RCF}$ could be proved directly: for $p(x) = x^{2n+1} + \sum_{i \leq 2n} a_i x^i$, by $\mathbf{FTA}_{\text{RCF}}$, there are $\{b_j, c_j\}_{j \leq n}$ such that $x \cdot p(x) = \prod_{j \leq n} (x^2 + b_j x + c_j)$. Since $\prod_{j \leq n} c_j = 0$, then for some j we have $c_j = 0$. Assume that $c_0 = 0$, then we have $x \cdot p(x) = x \cdot (x + b_0) \cdot \prod_{0 < j \leq n} (x^2 + b_j x + c_j)$ and so $p(-b_0) = 0$.

5 Conclusions and Open Problems

Continuous induction has been around in the literature over the last century (starting from [3] in 1919). We isolated three versions of it (Definitions 2.1, 2.3, 2.6, noting that all the other formats are equivalent to one of these three) and formalized them in first-order logic (Definitions 2.2, 2.5, 2.7). We showed that two versions of it are equivalent to each other (Theorem 2.8), while the third one (and the oldest one) is not (Theorem 2.9). Actually, those two strong versions are equivalent to the completeness of an ordered field, but the third one is equivalent to the Archimedean property (Theorem 3.8) and not with the completeness axiom. We noted that the first-order formulations of those two strong continuous induction schemes can completely axiomatize the real closed ordered fields (cf. Theorem 3.6). For this theory, we collected some axiomatizations (Definitions 3.1, 3.2, 3.4, 4.1, 4.2) which are shown to be equivalent to one another (Theorems 3.3, 3.5, 3.6) and added one; $\mathbf{FTA}_{\text{RCF}}$ a formalization of the fundamental theorem of algebra (Definition 4.10). The following diagram summarizes some of our new and old results (where the double lined arrows are proved here and the dotted arrows follow from Theorem 5.1 proved in the Appendix):



The completeness of **IVT** (+**OF**) is proved in the Appendix (Theorem 5.1); which, as was noted in Sect. 4, implies the equivalence of **D-Inf**(\mathcal{L}_{OF}) and **FTA**_{RCF} with **IVT**, as well.

1. We leave open the existence of a nice and neat first-order proof of any implication of the form **IVT** $\implies \Theta$, where Θ is either **FTA**_{RCF} or any of **D-Inf**, **D-Sup**, **D-Cut**, **DCI**, **DCI**₂ over \mathcal{L}_{OF} . Our intent by iterating some theorems from classical analysis was to put emphasis on their first-order formalizability. Indeed, being formalizable in first-order logic is not a trivial matter; all our proofs (except that of Theorem 3.8) were first-order.
2. The second open problem is a nice and neat first-order proof of **RCF** \implies **FTA**_{RCF} (closing the gap in the diagram). As we noted earlier, there are some second-order proofs for this (e.g., the proofs three and four for the fundamental theorem of algebra in [7]). By Gödel’s Completeness Theorem (for first-order logic) there should exist such proofs; but what are they? As a matter of fact, any such proof will be a beautiful proof of the fundamental theorem of algebra, which is completely real analytic (not referring to complex numbers) and it will be a *first-order proof* of the theorem, for the first time.
3. One research area which is wide open to explore is the formalization of the other known equivalent axiomatizations of the complete ordered fields in first-order logic, and seeing whether any of them can completely axiomatize the theory of real closed fields (over **OF**). Or more generally, is the theory **OF** + Θ , where Θ is a first-order formalization of any of the 72 completeness axioms in [5], equivalent to **RCF**? This question is not easy, since for example neither **OF** + **Rolle** nor **OF** + **MVT** is equivalent to **RCF** (see [2,13]), but e.g., **OF** + **DCI**(\mathcal{L}_{OF}) is equivalent to **RCF**. As we saw above, **OF** + Θ axiomatizes the real closed fields if and only if **OF** + $\Theta \vdash$ **IVT**. Let us note that **FTA**_{RCF} is not a first-order formalization of any of the completeness axioms in [5].

Acknowledgements The authors are very much grateful to the anonymous referee of the Bulletin for carefully reading the article and for many helpful suggestions which improved the results and made the paper much more readable.

Appendix

Here, we give a proof of Tarski’s theorem on the completeness of the theory of real closed (ordered) fields; this is a slightly modified proof given in [11].

Theorem 5.1 (*IVT is Complete*) *The theory **OF** + **IVT** is complete.*

Proof We will employ the method of quantifier elimination; i.e., we prove that every formula in the language of ordered fields is equivalent to a quantifier-free formula with

the same free variables, over the theory **OF+IVT**. Since this theory can decide (prove or refute) quantifier-free sentences, then the quantifier elimination theorem will show that **OF+IVT** can decide every sentence in its language; i.e., either proves it or proves its negation.

For that purpose, it suffices to show that every formula of the form $\exists x \theta(x)$, where θ is the conjunction of some atomic or negated-atomic formulas, is equivalent (in **OF+IVT**) to a quantifier-free formula (with the same free variables). To see this, suppose that every such formula is equivalent to a quantifier-free formula. Then by induction on the complexity of a formula ψ , one can show that ψ is equivalent to a quantifier-free formula: the case of atomic formulas and the propositional connectives $\{\neg, \wedge, \vee, \rightarrow\}$ are trivial, and the case of \forall can be reduced to that of \exists by $\forall x \psi(x) \equiv \neg \exists x \neg \psi(x)$. It remains to show the equivalence of $\exists x \psi(x)$ with a quantifier-free formula, where ψ is a quantifier-free formula. One can write ψ (equivalently) in disjunctive normal form $\bigvee_i \theta_i$ where each θ_i is a conjunction of some atomic or negated-atomic formulas. So, $\exists x \psi(x) \equiv \exists x \bigvee_i \theta_i \equiv \bigvee_i \exists x \theta_i$, and by the induction hypothesis each of $\exists x \theta_i$ is equivalent to some quantifier-free formula; thus the whole formula is so.

Since every term in the language $\{+, 0, -, \times, 1\}$ with the free variable x is a polynomial of x whose coefficients are some x -free terms, then every atomic formula with x is equivalent to $p(x) = 0$ or $q(x) > 0$ for some polynomials p, q . Noting that the negation sign \neg can be eliminated by $[p(x) \neq 0] \equiv [p(x) > 0 \vee p(x) < 0]$ and $[q(x) \neq 0] \equiv [q(x) = 0 \vee q(x) < 0]$, we can consider atomic formulas only. So, we consider the formulas of the following form for some polynomials $\{p_i(x)\}_{i < \ell}$ and $\{q_j(x)\}_{j < n}$: $\exists x [\bigwedge_{i < \ell} p_i(x) = 0 \wedge \bigwedge_{j < n} q_j(x) > 0]$. Finally, it suffices to consider the bounded formulas of the form $\exists x \in]a, b[: \psi(x)$ since every formula $\exists x \psi(x)$ is equivalent to $\exists x < -1 \psi(x) \vee \psi(-1) \vee \exists x \in]-1, 1[: \psi(x) \vee \psi(1) \vee \exists x > 1 \psi(x)$, and also we have $\exists x < -1 \psi(x) \equiv \exists y \in]0, 1[: \psi(-y^{-1})$ and $\exists x > 1 \psi(x) \equiv \exists y \in]0, 1[: \psi(y^{-1})$. Let us note that for any polynomials $p(x) = \sum_{i \leq m} a_i x^i$ and $q(x) = \sum_{j \leq l} b_j x^j$ and any $y > 0$ we have

$$\begin{cases} p(y^{-1}) = 0 \Leftrightarrow \sum_{i \leq m} a_i y^{-i} = 0 \Leftrightarrow \sum_{i \leq m} a_i y^{m-i} = 0 \Leftrightarrow \sum_{j \leq m} a_{m-j} y^j = 0, & \text{and} \\ q(y^{-1}) > 0 \Leftrightarrow \sum_{j \leq l} b_j y^{-j} > 0 \Leftrightarrow \sum_{j \leq l} b_j y^{l-j} > 0 \Leftrightarrow \sum_{i \leq l} a_{l-i} y^i > 0; \end{cases}$$

and so any atomic formula $\theta(y^{-1})$ (and also $\theta(-y^{-1})$), for $y > 0$, is equivalent to another atomic formula $\eta(y)$.

Whence, we show the equivalence of all the formulas in the following form with a quantifier-free formula:

$$(\dagger) \quad \varphi(a, b) = \exists x \in]a, b[: \bigwedge_{i < \ell} p_i(x) = 0 \wedge \bigwedge_{j < n} q_j(x) > 0.$$

This will be proved by induction on the degree of a formula which we define as follows:

- for a term $p(x) = \sum_{i \leq m} a_i x^i$, let $\deg_x p = m$;
- for atomic formulas $p(x) = 0$ or $q(x) > 0$, let $\deg_x(p(x) = 0) = \deg_x p$ and $\deg_x(q(x) > 0) = 1 + \deg_x q$;

– finally, the deg_x of a formula is the maximum of the deg_x 's of its atomic sub-formulas.

What we prove is:

(★) for any formula $\varphi(a, b)$ as (†) above, there are some formulas $\{\Phi_k(y), \Psi_k(y)\}_{k < m}$ such that $\mathbf{OF} + \mathbf{IVT} + a < b \vdash \varphi(a, b) \leftrightarrow \bigvee_{k < m} [\Phi_k(a) \wedge \Psi_k(b)]$, and moreover the deg_y of $\{\Phi_k(y), \Psi_k(y)\}$'s are less than $\text{deg}_x(\varphi(a, b))$.

This will be shown by induction on $\bar{h} = \text{deg}_x(\varphi(a, b))$; let us note that here a, b are treated as (new) variables. If $\bar{h} = 0$, then x appears only superficially in $\varphi(a, b)$ and so it is equivalent to a quantifier-free formula. Now, suppose that we have the desired conclusion (★) for all the formulas with deg_x less than \bar{h} .

(1) First, we consider the case of $\ell = 0$; i.e., the formulas that are in the following form:

$$(†) \quad \varphi(a, b) = \exists x \in]a, b[: \bigwedge_{j < n} q_j(x) > 0.$$

By Lemma 4.7, q_j 's are positive at a point in $]a, b[$ if and only if they are positive in some sub-interval $]u, v[\subseteq]a, b[$. So, we have $\varphi(a, b) \equiv F(a, b) \vee \bigvee_{i < n} G_i(a, b) \vee \bigvee_{i, j < n} H_{i, j}(a, b)$, where $F, \{G_i\}_{i < n}$ and $\{H_{i, j}\}_{i, j < n}$ are as follows:

$$\begin{aligned} F(a, b) &= \forall x \in]a, b[: \bigwedge_{j < n} q_j(x) > 0, \\ G_i(a, b) &= [\exists u \in]a, b[: q_i(u) = 0 \wedge F(a, u)] \vee [\exists v \in]a, b[: q_i(v) = 0 \wedge F(v, b)], \\ &\text{and} \\ H_{i, j}(a, b) &= \exists u, v \in]a, b[: u < v \wedge q_i(u) = 0 \wedge q_j(v) = 0 \wedge F(u, v). \end{aligned}$$

The formula $F(a, b)$ is equivalent to $\bigwedge_{j < n} q_j(a) \geq 0 \wedge \bigwedge_{j < n} \neg \exists x \in]a, b[: q_j(x) = 0$, whose deg_x is less than \bar{h} , and so by the induction hypothesis (★) is equivalent to the formula $\bigvee_{k < m} [\Phi_k(a) \wedge \Psi_k(b)]$ for some x -free formulas $\{\Phi_k(y), \Psi_k(y)\}_{k < m}$ whose deg_y are less than \bar{h} . So, for any $i < n$, $G_i(a, b)$ is equivalent to the following formula:

$$\begin{aligned} &\bigvee_{k < m} [(\Phi_k(a) \wedge \exists u \in]a, b[: [q_i(u) = 0 \wedge \Psi_k(u)]) \\ &\vee (\exists v \in]a, b[: [q_i(v) = 0 \wedge \Phi_k(v)] \wedge \Psi_k(b))]. \end{aligned}$$

Since the deg_u of $q_i(u) = 0 \wedge \Psi_k(u)$ and the deg_v of $q_i(v) = 0 \wedge \Phi_k(v)$ are less than \bar{h} , the induction hypothesis (★) applies to all G_i 's. Finally, for $H_{i, j}(a, b)$, we note that it is equivalent to the disjunction (over $k < m$) of the following formulas:

$$\exists u \in]a, b[: (q_i(u) = 0 \wedge \Phi_k(u) \wedge \exists v \in]u, b[: [q_j(v) = 0 \wedge \Psi_k(v)]).$$

Now, the formula $q_j(v) = 0 \wedge \Psi_k(v)$ has deg_v less than \bar{h} ; so by the induction hypothesis (★), there are formulas $\{\Theta_{j, k, l}(z), \Upsilon_{j, k, l}(z)\}_{l < l}$ with deg_z less than \bar{h} such that the formula $\exists v \in]u, b[: [q_j(v) = 0 \wedge \Psi_k(v)]$ is equivalent to $\bigvee_{l < l} [\Theta_{j, k, l}(u) \wedge \Upsilon_{j, k, l}(b)]$. Thus, $H_{i, j}(a, b)$ is equivalent to the disjunction of $\exists u \in]a, b[: (q_i(u) = 0 \wedge \Phi_k(u) \wedge$

$\Theta_{j,k,\iota}(u) \wedge \Upsilon_{i,j,\iota}(b)$, for $k < m$, $\iota < l$, to which the induction hypothesis (\star) apply, since the \deg_u of the all the formulas $q_i(u) = 0 \wedge \Phi_k(u) \wedge \Theta_{j,k,\iota}(u)$ are less than \bar{h} .

(2) Second, we consider the case of $\ell > 0$; let us note that we can assume $\ell = 1$ since we have $\bigwedge_{i < \ell} \alpha_i = 0 \iff \sum_{i < \ell} \alpha_i^2 = 0$. So, we may replace all the $p_i(x)$'s with a single polynomial $\sum_{i < \ell} p_i^2(x)$; but this will increase the \deg_x of the resulted formula. There is another way of reducing ℓ (the number of polynomials p_i 's) without increasing the \deg_x of the formula:

(I) For polynomials $p(x) = \alpha x^d + \sum_{i < d} a_i x^i$ and $q(x) = \beta x^e + \sum_{j < e} b_j x^j$ assume that $d \geq e$. Put $r(x) = q(x) - \beta x^e$ and $s(x) = \beta p(x) - \alpha x^{d-e} q(x)$; then we have $[p(u) = q(u) = 0] \iff [\beta = 0 \wedge p(u) = r(u) = 0] \vee [\beta \neq 0 \wedge s(u) = q(u) = 0]$.

Continuing this way, at least one of the two polynomials will disappear and we will be left with at most one polynomial, and the \deg_x of the last formula will be non-greater than the \deg_x of the first formula.

So, we can safely assume that $\ell = 1$; thus $\varphi(a, b) = \exists x \in]a, b[: p(x) = 0 \wedge \bigwedge_{j < n} q_j(x) > 0$ and $\deg_x(\varphi(a, b)) = \bar{h}$. We can still transform the formula to an equivalent one in which we have $\deg_x q_j < \deg_x p$ for all $j < n$:

(II) If, say, $\deg_x q_1 \geq \deg_x p$, then write $p(x) = \alpha x^d + \sum_{i < d} a_i x^i$ and $q_1(x) = \beta x^e + \sum_{j < e} b_j x^j$ with $d \leq e$. Put $r(x) = p(x) - \alpha x^d$ and $s(x) = \alpha^2 q_1(x) - \alpha \beta x^{e-d} p(x)$; then we have $[p(u) = 0 \wedge q_1(u) > 0] \iff [\alpha = 0 \wedge r(u) = 0 \wedge q_1(u) > 0] \vee [\alpha \neq 0 \wedge p(u) = 0 \wedge s(u) > 0]$. Continuing this way, either the equality ($p(x) = 0$) will disappear (and so we will have the first case) or the degree of the inequality ($q_1(x) > 0$) will be non-greater than the degree of the equality, $\deg_x(p(x) = 0)$.

So, assume that in the formula

$$(\dagger) \quad \varphi(a, b) = \exists x \in]a, b[: p(x) = 0 \wedge \bigwedge_{j < n} q_j(x) > 0,$$

we have that $\deg_x(\varphi(a, b)) = \bar{h} = \deg_x p$ and also $\bigwedge_{j < n} \deg_x q_j < \bar{h}$. Now, the formula $\varphi(a, b)$ is equivalent to $\varphi_1(a, b) \vee \varphi_2(a, b) \vee \varphi_3(a, b)$, where

$$\begin{aligned} \varphi_1(a, b) &= \exists x \in]a, b[: p(x) = 0 \wedge p'(x) = 0 \wedge \bigwedge_{j < n} q_j(x) > 0, \\ \varphi_2(a, b) &= \exists x \in]a, b[: p(x) = 0 \wedge p'(x) > 0 \wedge \bigwedge_{j < n} q_j(x) > 0, \text{ and} \\ \varphi_3(a, b) &= \exists x \in]a, b[: p(x) = 0 \wedge p'(x) < 0 \wedge \bigwedge_{j < n} q_j(x) > 0. \end{aligned}$$

Since, $\deg_x p' < \deg_x p$ (see Definition 4.4), then by applying (I) to φ_1 we get an equivalent formula with \deg_x less than \bar{h} and then can apply the induction hypothesis (\star) . For φ_2 , we note that, by **IVT** and Theorem 4.6, all the q_j s and also p' are strictly positive at some point in $]a, b[$ in which p vanishes, if and only if all the q_j s and p' are strictly positive on some open sub-interval $]u, v[\subseteq]a, b[$ such that (p is monotonically increasing and so) $p(u) < 0 < p(v)$. Whence, $\varphi_2(a, b)$ is equivalent to the disjunction of the following formulas:

- (i) $p(a) < 0 < p(b) \wedge F(a, b)$,
- (ii) $\bigvee_{i < n} \left([p(a) < 0 \wedge \exists u \in]a, b[: (q_i(u) = 0 \wedge p(u) > 0 \wedge F(a, u))] \vee [p(b) > 0 \wedge \exists v \in]a, b[: (q_i(v) = 0 \wedge p(v) < 0 \wedge F(v, b))] \right)$, and
- (iii) $\bigvee_{i, j < n} \left(\exists u, v \in]a, b[: [q_i(u) = 0 \wedge q_j(v) = 0 \wedge p(u) < 0 < p(v) \wedge F(u, v)] \right)$.

The formula (i) has been treated before (it is equivalent to a formula with deg_x less than \bar{h}). The formulas (ii) and (iii) can be equivalently transformed to formulas with deg_x less than \bar{h} by (I) and (II) above. So, the whole formula φ_2 , and very similarly φ_3 , can be written in equivalent forms in such a way that the induction hypothesis (\star) applies to them. \square

References

1. Basu, S., Pollack, R., Roy, M.-F.: Algorithms in Real Algebraic Geometry, 2nd edn. Springer, Berlin (2006). ISBN:9783540330981
2. Brown, R., Craven, T.C., Pelling, M.J.: Ordered fields satisfying Rolle's theorem. III. J. Math. **30**(1), 66–78 (1986). <https://projecteuclid.org/euclid.ijm/1256044752>
3. Chao, Y.R.: A note on "Continuous mathematical induction". Bull. Am. Math. Soc. **26**(1) 17–18 (1919). <https://projecteuclid.org/euclid.bams/1183425067>
4. Clark, P.L.: The instructor's guide to real induction. Expository preprint (2017). Available on the net at: http://alpha.math.uga.edu/~pete/instructors_guide_2017.pdf. Accessed 15 May 2019
5. Deveau, M., Teismann, H.: 72+42: Characterizations of the completeness and Archimedean properties of ordered fields. Real Anal. Exch. **39**(2), 261–304 (2014). <https://doi.org/10.14321/realanalexch.39.2.0261>
6. Epstein, R.L.: Classical Mathematical Logic: The Semantic Foundations of Logic. Princeton University Press, Princeton (2006). (corr. 2011). ISBN:9780691123004
7. Fine, B., Rosenberger, G.: The Fundamental Theorem of Algebra. Springer, Berlin (1997). (softcover 2012). ISBN:9781461273431
8. Hathaway, D.: Using continuity induction. Coll. Math. J. (Classr. Capsul.) **42**(3), 229–231 (2011). <https://doi.org/10.4169/college.math.j.42.3.229>
9. Jingzhong, Z.: The induction on a continuous variable. IAEA-UNESCO, ICTP Report #IC/89/157 (1989). <http://streaming.ictp.it/preprints/P/89/157.pdf>. Accessed 15 May 2019
10. Kalantari, I.: Induction over the continuum. In: Friend, M., Goethe, N.B., Harizanov, V.S. (eds.) Induction, Algorithmic Learning Theory, and Philosophy, pp. 145–154. Springer, Berlin (2007). https://doi.org/10.1007/978-1-4020-6127-1_5
11. Kreisel, G., Krivine, J.-L.: Elements of Mathematical Logic: Model Theory. North-Holland, Amsterdam (1971). ISBN:9780720422658
12. Mashkouri, M.: Sākhtāre A'dād, Mabāniye Riyāziāt. Structure of Numbers, Foundations of Mathematics. Isfahan University of Technology Publication Center, Isfahan (2004). ISBN:9648476012 (in Farsi). <https://publication.iut.ac.ir/book/89>
13. Pelling, M.J.: Solution of advanced problem no. 5861 (proposed by Michael Slater). Am. Math. Mon. **88**(2), 150–152 (1981). <https://www.jstor.org/stable/2321145>
14. Spivak, M.: Calculus, Publish or Perish, Houston, 4th edn (2008). ISBN:9780914098911

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.